

References

1. Abstracts of the the american society for neurochemistry 37th annual meeting, portland, oregon, USA, 11-15 march 2006. *J Neurochem* 2006 Mar;96 Suppl 1:1-150.
2. *Annals of Neurology*(5):- 625.
3. Aggarwal S, Jinda W, Limwongse C, Atchaneeyasakul LO, Phadke SR. Run-on mutation in the PAX6 gene and chorioretinal degeneration in autosomal dominant aniridia. *Mol Vis* 2011;17:1305-9.
4. Akiyama M, Titeux M, Sakai K, McMillan JR, Tonasso L, Calvas P, Jossic F, Hovnanian A, Shimizu H. DNA-based prenatal diagnosis of harlequin ichthyosis and characterization of ABCA12 mutation consequences. *J Invest Dermatol* 2007 Mar;127(3):568-73.
5. Akiyama M, Titeux M, Sakai K, McMillan JR, Tonasso L, Calvas P, Jossic F, Hovnanian A, Shimizu H. DNA-based prenatal diagnosis of harlequin ichthyosis and characterization of ABCA12 mutation consequences RID C-6657-2009. *J Invest Dermatol* 2007 MAR;127(3):568-73.
6. Alcantara-Ortigoza MA, Belmont-Martinez L, Vela-Amieva M, Gonzalez-Del Angel A. Analysis of the CTNS gene in nephropathic cystinosis mexican patients: Report of four novel mutations and identification of a false positive 57-kb deletion genotype with LDM-2/exon 4 multiplex PCR assay. *Genet Test* 2008 Sep;12(3):409-14.
7. Allikmets R, Wasserman WW, Hutchinson A, Smallwood P, Nathans J, Rogan PK, Schneider TD, Dean M. Organization of the ABCR gene: Analysis of promoter and splice junction sequences. *Gene* 1998 Jul 17;215(1):111-22.
8. Anczukow O, Buisson M, Salles MJ, Triboulet S, Longy M, Lidereau R, Sinilnikova OM, Mazoyer S. Unclassified variants identified in BRCA1 exon 11: Consequences on splicing. *Genes Chromosomes Cancer* 2008 May;47(5):418-26.
9. Aoyama Y, Ozer I, Demirkol M, Ebara T, Murase T, Podskarbi T, Shin YS, Gokcay G, Okubo M. Molecular features of 23 patients with glycogen storage disease type III in turkey: A novel mutation p.R1147G associated with isolated glucosidase deficiency, along with 9 AGL mutations. *J Hum Genet* 2009 NOV;54(11):681-6.
10. Arita K, Wessagowit V, Inamadar AC, Palit A, Fassihi H, Lai-Cheong JE, Poureyyron C, South AP, McGrath JA. Unusual molecular findings in kindler syndrome. *Br J Dermatol* 2007 Dec;157(6):1252-6.
11. Atwood CS, inventor. Methods of assessing risk of alzheimer's disease in a patient. United States patent 2009.
12. Bacci C, Sestini R, Provenzano A, Paganini I, Mancini I, Porfirio B, Vivarelli R, Genuardi M, Papi L. Schwannomatosis associated with multiple meningiomas due to a familial SMARCB1 mutation. *Neurogenetics* 2010 Feb;11(1):73-80.
13. Baralle D, Lucassen A, Buratti E. Missed threads. the impact of pre-mRNA splicing defects on clinical practice. *EMBO Rep* 2009 Aug;10(8):810-6.
14. Beetz C, Schule R, Deconinck T, Tran-Viet KN, Zhu H, Kremer BP, Frints SG, van Zelst-Stams WA, Byrne P, Otto S, et al. REEP1 mutation spectrum and genotype/phenotype correlation in hereditary spastic paraplegia type 31. *Brain* 2008 Apr;131(Pt 4):1078-86.

15. Ben Selma Z, Yilmaz S, Schischmanoff PO, Blom A, Ozogul C, Laroche L, Caux F. A novel S115G mutation of CGI-58 in a turkish patient with dorfman-chanarin syndrome. *J Invest Dermatol* 2007 Sep;127(9):2273-6.
16. Bertola F, Filocamo M, Casati G, Mort M, Rosano C, Tylki-Szymanska A, Tuysuz B, Gabrielli O, Grossi S, Scarpa M, et al. IDUA mutational profiling of a cohort of 102 european patients with mucopolysaccharidosis type I: Identification and characterization of 35 novel alpha-L-iduronidase (IDUA) alleles. *Hum Mutat* 2011 Jun;32(6):E2189-210.
17. Bi C, Rogan P. Determining thresholds for binding site sequence models using information theory. ; 2005. PT: B; CT: 8th Joint Conference on Information Sciences (JCIS 2005); CY: JUL 21-26, 2005; CL: Salt Lake City, UT; TC: 1; UT: WOS:000233670802026.
18. Bloethner S, Mould A, Stark M, Hayward NK. Identification of ARHGEF17, DENND2D, FGFR3, and RB1 mutations in melanoma by inhibition of nonsense-mediated mRNA decay. *Genes Chromosomes Cancer* 2008 Dec;47(12):1076-85.
19. Bocchi L, Pisciotta L, Fasano T, Candini C, Puntoni MR, Sampietro T, Bertolini S, Calandra S. Multiple abnormally spliced ABCA1 mRNAs caused by a novel splice site mutation of ABCA1 gene in a patient with tangier disease. *Clin Chim Acta* 2010 Apr 2;411(7-8):524-30.
20. Bogaerts V, Nuytemans K, Reumers J, Pals P, Engelborghs S, Pickut B, Corsmit E, Peeters K, Schymkowitz J, De Deyn PP, et al. Genetic variability in the mitochondrial serine protease HTRA2 contributes to risk for parkinson disease. *Hum Mutat* 2008 Jun;29(6):832-40.
21. Bonnet-Dupeyron MN, Combes P, Santander P, Cailloux F, Boespflug-Tanguy O, Vaur-Barriere C. PLP1 splicing abnormalities identified in pelizaeus-merzbacher disease and SPG2 fibroblasts are associated with different types of mutations. *Hum Mutat* 2008 Aug;29(8):1028-36.
22. Borroni B, Archetti S, Alberici A, Agosti C, Gennarelli M, Bigni B, Bonvicini C, Ferrari M, Bellelli G, Galimberti D, et al. Progranulin genetic variations in frontotemporal lobar degeneration: Evidence for low mutation frequency in an italian clinical series. *Neurogenetics* 2008 Jul;9(3):197-205.
23. Botta E, Nardo T, Orioli D, Guglielmino R, Ricotti R, Bondanza S, Benedicenti F, Zambruno G, Stefanini M. Genotype-phenotype relationships in trichothiodystrophy patients with novel splicing mutations in the XPD gene. *Hum Mutat* 2009 Mar;30(3):438-45.
24. Brockmoller J, Tzvetkov MV. Pharmacogenetics: Data, concepts and tools to improve drug discovery and drug treatment. *Eur J Clin Pharmacol* 2008 Feb;64(2):133-57.
25. Broer S, Bailey CG, Kowalczyk S, Ng C, Vanslambrouck JM, Rodgers H, Auray-Blais C, Cavanaugh JA, Broer A, Rasko JE. Iminoglycinuria and hyperglycinuria are discrete human phenotypes resulting from complex mutations in proline and glycine transporters. *J Clin Invest* 2008 Dec;118(12):3881-92.
26. Buratti E, Baralle M, Baralle FE. Defective splicing, disease and therapy: Searching for master checkpoints in exon definition. *Nucleic Acids Res* 2006 Jul 19;34(12):3494-510.
27. Cabral RM, Liu L, Hogan C, Dopping-Hepenstal PJ, Winik BC, Asial RA, Dobson R, Mein CA, Baselaga PA, Mellerio JE, et al. Homozygous mutations in the 5' region of the JUP gene

- result in cutaneous disease but normal heart development in children. *J Invest Dermatol* 2010 Jun;130(6):1543-50.
28. Calandra S, Tarugi P, Bertolini S. Altered mRNA splicing in lipoprotein disorders. *Curr Opin Lipidol* 2011 Apr;22(2):93-9.
 29. Caridi G, Dagnino M, Dalgic B, Egritas O, Sancak B, Campagnoli M, Dolcini L, Galliano M, Minchiotti L. Analbuminemia zonguldak: Case report and mutational analysis. *Clin Biochem* 2008 Mar;41(4-5):288-91.
 30. Castaman G, Giacomelli SH, Mancuso ME, D'Andrea G, Santacroce R, Sanna S, Santagostino E, Mannucci PM, Goodeve A, Rodeghiero F. Deep intronic variations may cause mild hemophilia A. *J Thromb Haemost* 2011 Aug;9(8):1541-8.
 31. Castiglia D, Zambruno G. Mutation mechanisms. *Dermatol Clin* 2010 Jan;28(1):17-22.
 32. Caux-Moncoutier V, Pages-Berhouet S, Michaux D, Asselain B, Castera L, De Pauw A, Buecher B, Gauthier-Villars M, Stoppa-Lyonnet D, Houdayer C. Impact of BRCA1 and BRCA2 variants on splicing: Clues from an allelic imbalance study. *Eur J Hum Genet* 2009 Nov;17(11):1471-80.
 33. Cefalu AB, Noto D, Magnolo L, Pinotti E, Gomaraschi M, Martini S, Vigna GB, Calabresi L, Tarugi P, Aversa MR. Novel mutations of CETP gene in Italian subjects with hyperalphalipoproteinemia. *Atherosclerosis* 2009 May;204(1):202-7.
 34. Chen L, Qin S, Xie J, Tang J, Yang L, Shen W, Zhao X, Du J, He G, Feng G, et al. Genetic polymorphism analysis of CYP2C19 in Chinese Han populations from different geographic areas of mainland China. *Pharmacogenomics* 2008 Jun;9(6):691-702.
 35. Chen LJ, Tam PO, Tham CC, Liang XY, Chiang SW, Canlas O, Ritch R, Rhee DJ, Pang CP. Evaluation of SPARC as a candidate gene of juvenile-onset primary open-angle glaucoma by mutation and copy number analyses. *Mol Vis* 2010 Oct 8;16:2016-25.
 36. Chen LJ, Tam POS, Tham CCY, Liang XY, Chiang SWY, Canlas O, Ritch R, Rhee DJ, Pang CP. Evaluation of SPARC as a candidate gene of juvenile-onset primary open-angle glaucoma by mutation and copy number analyses. *Molecular Vision* 2010 OCT 8;16(215-19):2016-25.
 37. Clark GR, Crowe P, Muszynska D, O'Prey D, O'Neill J, Alexander S, Willoughby CE, McKay GJ, Silvestri G, Simpson DA. Development of a diagnostic genetic test for simplex and autosomal recessive retinitis pigmentosa. *Ophthalmology* 2010 Nov;117(11):2169,77.e3.
 38. Concolino P, Vendittelli F, Mello E, Minucci A, Carrozza C, Rossodivita A, Giardina B, Zuppi C, Capoluongo E. Functional analysis of two rare CYP21A2 mutations detected in Italian patients with a mildest form of congenital adrenal hyperplasia. *Clin Endocrinol (Oxf)* 2009 Oct;71(4):470-6.
 39. Covaciu C, Grosso F, Pisaneschi E, Zambruno G, Gregersen PA, Sommerlund M, Hertz JM, Castiglia D. A founder synonymous COL7A1 mutation in three Danish families with dominant dystrophic epidermolysis bullosa pruriginosa identifies exonic regulatory sequences required for exon 87 splicing. *Br J Dermatol* 2011 Sep;165(3):678-82.
 40. Cox DG, Crusius JB, Peeters PH, Bueno-de-Mesquita HB, Pena AS, Canzian F. Haplotype of prostaglandin synthase 2/cyclooxygenase 2 is involved in the susceptibility to inflammatory bowel disease. *World J Gastroenterol* 2005 Oct 14;11(38):6003-8.

41. Cruchaga C, Fernandez-Seara MA, Seijo-Martinez M, Samaranch L, Lorenzo E, Hinrichs A, Irigoyen J, Maestro C, Prieto E, Marti-Clement JM, et al. Cortical atrophy and language network reorganization associated with a novel progranulin mutation. *Cereb Cortex* 2009 Aug;19(8):1751-60.
42. Dash DP, George S, O'Prey D, Burns D, Nabili S, Donnelly U, Hughes AE, Silvestri G, Jackson J, Frazer D, et al. Mutational screening of VSX1 in keratoconus patients from the european population. *Eye (Lond)* 2010 Jun;24(6):1085-92.
43. IDDM2 locus: 5' noncoding intron I splicing and translational efficiency effects of INS - 23HphI - more than a tag for the INS promoter VNTR. [Internet]; c2006. Available from: abstracts/hgvs.org/Helsinki/Presentations/Day.ppt.
44. Desmet FO, Hamroun D, Lalande M, Collod-Beroud G, Claustres M, Beroud C. Human splicing finder: An online bioinformatics tool to predict splicing signals. *Nucleic Acids Res* 2009 May;37(9):e67.
45. Di Leo E, Magnolo L, Lancellotti S, Croce L, Visintin L, Tiribelli C, Bertolini S, Calandra S, Tarugi P. Abnormal apolipoprotein B pre-mRNA splicing in patients with familial hypobetalipoproteinaemia. *J Med Genet* 2007 Mar;44(3):219-24.
46. Di Leo E, Magnolo L, Pinotti E, Martini S, Cortella I, Vitturi N, Rabacchi C, Wunsch A, Pucci F, Bertolini S, et al. Functional analysis of two novel splice site mutations of APOB gene in familial hypobetalipoproteinemia. *Mol Genet Metab* 2009 Feb;96(2):66-72.
47. Di Leo E, Panico F, Tarugi P, Battisti C, Federico A, Calandra S. A point mutation in the lariat branch point of intron 6 of NPC1 as the cause of abnormal pre-mRNA splicing in niemann-pick type C disease. *Hum Mutat* 2004 Nov;24(5):440.
48. Dinakarbandian D, Raheja V, Mehta S, Schuetz E, Rogan P. Tandem machine learning for the identification of genes regulated by transcription factors. *BMC Bioinformatics* 2005 AUG 22;6:204.
49. Douglas DA, Zhong H, Ro JY, Oddoux C, Berger AD, Pincus MR, Satagopan JM, Gerald WL, Scher HI, Lee P, et al. Novel mutations of epidermal growth factor receptor in localized prostate cancer. *Front Biosci* 2006 Sep 1;11:2518-25.
50. Drogemuller C, Philipp U, Haase B, Gunzel-Apel AR, Leeb T. A noncoding melanophilin gene (MLPH) SNP at the splice donor of exon 1 represents a candidate causal mutation for coat color dilution in dogs. *J Hered* 2007;98(5):468-73.
51. Dua-Awereh MB, Shimomura Y, Kraemer L, Wajid M, Christiano AM. Mutations in the desmoglein 1 gene in five pakistani families with striate palmoplantar keratoderma. *J Dermatol Sci* 2009 Mar;53(3):192-7.
52. Dutrannoy V, Demuth I, Baumann U, Schindler D, Konrat K, Neitzel H, Gillissen-Kaesbach G, Radszewski J, Rothe S, Schellenberger MT, et al. Clinical variability and novel mutations in the NHEJ1 gene in patients with a nijmegen breakage syndrome-like phenotype. *Hum Mutat* 2010 Sep;31(9):1059-68.
53. Eckl KM, de Juanes S, Kurtenbach J, Natebus M, Lugassy J, Oji V, Traupe H, Preil ML, Martinez F, Smolle J, et al. Molecular analysis of 250 patients with autosomal recessive congenital ichthyosis: Evidence for mutation hotspots in ALOXE3 and allelic heterogeneity in ALOX12B. *J Invest Dermatol* 2009 Jun;129(6):1421-8.

54. Ellis JR, Jr, Heinrich B, Mautner VF, Kluwe L. Effects of splicing mutations on NF2-transcripts: Transcript analysis and information theoretic predictions. *Genes Chromosomes Cancer* 2011 Aug;50(8):571-84.
55. ElSharawy A, Hundrieser B, Brosch M, Wittig M, Huse K, Platzer M, Becker A, Simon M, Rosenstiel P, Schreiber S, et al. Systematic evaluation of the effect of common SNPs on pre-mRNA splicing. *Hum Mutat* 2009 Apr;30(4):625-32.
56. Fang S, Guo X, Jia X, Xiao X, Li S, Zhang Q. Novel GPR143 mutations and clinical characteristics in six chinese families with X-linked ocular albinism. *Mol Vis* 2008;14:1974-82.
57. Fasano T, Bocchi L, Pisciotta L, Bertolini S, Calandra S. Denaturing high-performance liquid chromatography in the detection of ABCA1 gene mutations in familial HDL deficiency. *J Lipid Res* 2005 Apr;46(4):817-22.
58. Fasano T, Pisciotta L, Bocchi L, Guardamagna O, Assandro P, Rabacchi C, Zanoni P, Filocamo M, Bertolini S, Calandra S. Lysosomal lipase deficiency: Molecular characterization of eleven patients with wolman or cholesteryl ester storage disease. *Mol Genet Metab* 2011 Dec 17.
59. Fattal-Valevski A, DiMaio MS, Hisama FM, Hobson GM, Davis-Williams A, Garbern JY, Mahoney MJ, Kolodny EH, Pastores GM. Variable expression of a novel PLP1 mutation in members of a family with pelizaeus-merzbacher disease. *J Child Neurol* 2009 May;24(5):618-24.
60. Faz DB. Bases geneticas de la conducta / genetic bases of behavior. Barcelona, Catalonia, Spain: Edicions de la Universitat Oberta de Catalunya; 2009. .
61. Fornage M, Lee CR, Doris PA, Bray MS, Heiss G, Zeldin DC, Boerwinkle E. The soluble epoxide hydrolase gene harbors sequence variation associated with susceptibility to and protection from incident ischemic stroke. *Hum Mol Genet* 2005 Oct 1;14(19):2829-37.
62. Gaedigk A, Baker DW, Totah RA, Gaedigk R, Pearce RE, Vyhldal CA, Zeldin DC, Leeder JS. Variability of CYP2J2 expression in human fetal tissues. *J Pharmacol Exp Ther* 2006 Nov;319(2):523-32.
63. Gaedigk A, Bhatena A, Ndjountche L, Pearce RE, Abdel-Rahman SM, Alander SW, Bradford LD, Rogan PK, Leeder JS. Identification and characterization of novel sequence variations in the cytochrome P4502D6 (CYP2D6) gene in african americans. *Pharmacogenomics J* 2005;5(3):173-82.
65. Gaedigk A, Gaedigk R, Leeder JS. CYP2D7 splice variants in human liver and brain: Does CYP2D7 encode functional protein? *Biochem Biophys Res Commun* 2005 Nov 4;336(4):1241-50.
66. Gaedigk A, Leeder JS. Letter to the editor. *Clin Pharmacol Ther* 2006;80:558-560.
67. Gaedigk A, Ndjountche L, Leeder JS, Bradford LD. Limited association of the 2988g > a single nucleotide polymorphism with CYP2D641 in black subjects. *Clin Pharmacol Ther* 2005 Mar;77(3):228,30; author reply 230-1.
68. Gao S, Zhang N, Zhang L, Duan GY, Zhang T. The human variome project and its progress. *Yi Chuan* 2010 Nov;32(11):1105-13.

69. Garcia-Blanco M. Alternative splicing: Therapeutic target and tool. In: P. Jeanteur, editor. *Alternative splicing and disease*. Berlin, Germany: Springer; 2006. .
70. Garcia-Gonzalez MA, Jones JG, Allen SK, Palatucci CM, Batish SD, Seltzer WK, Lan Z, Allen E, Qian F, Lens XM, et al. Evaluating the clinical utility of a molecular genetic test for polycystic kidney disease. *Mol Genet Metab* 2007 Sep-Oct;92(1-2):160-7.
71. Gaweda-Walerych K, Safranow K, Maruszak A, Bialecka M, Klodowska-Duda G, Czyzewski K, Slawek J, Rudzinska M, Styczynska M, Opala G, et al. Mitochondrial transcription factor A variants and the risk of parkinson's disease. *Neurosci Lett* 2010 Jan 18;469(1):24-9.
73. Gerykov-Bujalkova M, Krivulcik T, Bartosova Z. Novel approaches in evaluation of pathogenicity of single-base exonic germline changes involving the mismatch repair genes MLH1 and MSH2 in diagnostics of lynch syndrome minireview. *Neoplasma* 2008;55(6):463-71.
74. Godefroid N, Riveira-Munoz E, Saint-Martin C, Nassogne MC, Dahan K, Devuyst O. A novel splicing mutation in SLC12A3 associated with gitelman syndrome and idiopathic intracranial hypertension. *Am J Kidney Dis* 2006 Nov;48(5):e73-9.
75. Gruber FX, Hjorth-Hansen H, Mikkola I, Stenke L, Johansen T. A novel bcr-abl splice isoform is associated with the L248V mutation in CML patients with acquired resistance to imatinib. *Leukemia* 2006 Nov;20(11):2057-60.
76. Hageman GS, Anderson DH, Johnson LV, Hancox LS, Taiber AJ, Hardisty LI, Hageman JL, Stockman HA, Borchardt JD, Gehrs KM, et al. A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc Natl Acad Sci U S A* 2005 May 17;102(20):7227-32.
77. Hamada T, Fukuda S, Sakaguchi S, Yasumoto S, Kim SC, Hashimoto T. Molecular and clinical characterization in japanese and korean patients with hailey-hailey disease: Six new mutations in the ATP2C1 gene. *J Dermatol Sci* 2008 Jul;51(1):31-6.
78. Hampson G, Konrad MA, Scoble J. Familial hypomagnesaemia with hypercalciuria and nephrocalcinosis (FHHNC): Compound heterozygous mutation in the claudin 16 (CLDN16) gene. *BMC Nephrol* 2008 Sep 24;9:12.
79. Hartmann L, Theiss S, Niederacher D, Schaal H. Diagnostics of pathogenic splicing mutations: Does bioinformatics cover all bases? *Front Biosci* 2008 May 1;13:3252-72.
80. Hellerud C, Adamowicz M, Jurkiewicz D, Taybert J, Kubalska J, Ciara E, Popowska E, Ellis JR, Lindstedt S, Pronicka E. Clinical heterogeneity and molecular findings in five polish patients with glycerol kinase deficiency: Investigation of two splice site mutations with computerized splice junction analysis and Xp21 gene-specific mRNA analysis. *Mol Genet Metab* 2003 Jul;79(3):149-59.
81. Henneman P, Schaap FG, Rensen PC, van Dijk KW, Smelt AH. Estrogen induced hypertriglyceridemia in an apolipoprotein AV deficient patient. *J Intern Med* 2008 Jan;263(1):107-8.
82. Henriksen AM, Tumer Z, Tommerup N, Tranebjaerg L, Larsen LA. Identification of a novel EYA1 splice-site mutation in a danish branchio-oto-renal syndrome family. *Genet Test* 2004 Winter;8(4):404-6.

83. Hiller M, Huse K, Szafranski K, Rosenstiel P, Schreiber S, Backofen R, Platzer M. Phylogenetically widespread alternative splicing at unusual GYNGYN donors. *Genome Biol* 2006;7(7):R65.
84. Hines RN, Koukouritaki SB, Poch MT, Stephens MC. Regulatory polymorphisms and their contribution to interindividual differences in the expression of enzymes influencing drug and toxicant disposition. *Drug Metab Rev* 2008;40(2):263-301.
85. Hobson GM, Huang Z, Sperle K, Sistermans E, Rogan PK, Garbern JY, Kolodny E, Naidu S, Cambi F. Splice-site contribution in alternative splicing of PLP1 and DM20: Molecular studies in oligodendrocytes. *Hum Mutat* 2006 Jan;27(1):69-77.
86. Hobson GM, Huang Z, Sperle K, Sistermans E, Rogan PK, Garbern JY, Kolodny E, Naidu S, Cambi F. Splice-site contribution in alternative splicing of PLP1 and DM20: Molecular studies in oligodendrocytes. *Hum Mutat* 2006 Jan;27(1):69-77.
87. Houdayer C, Dehainault C, Mattler C, Michaux D, Caux-Moncoutier V, Pages-Berhouet S, d'Enghien CD, Lauge A, Castera L, Gauthier-Villars M, et al. Evaluation of in silico splice tools for decision-making in molecular diagnosis. *Hum Mutat* 2008 Jul;29(7):975-82.
88. Hube F, Guo J, Chooniedass-Kothari S, Cooper C, Hamedani MK, Dibrov AA, Blanchard AA, Wang X, Deng G, Myal Y, et al. Alternative splicing of the first intron of the steroid receptor RNA activator (SRA) participates in the generation of coding and noncoding RNA isoforms in breast cancer cell lines. *DNA Cell Biol* 2006 Jul;25(7):418-28.
89. Inui H, Oh KS, Nadem C, Ueda T, Khan SG, Metin A, Gozukara E, Emmert S, Slor H, Busch DB, et al. Xeroderma pigmentosum-variant patients from america, europe, and asia. *J Invest Dermatol* 2008 Aug;128(8):2055-68.
90. Johnson DS. Study of a possible genetic cause of CHARGE association. 2010.
91. Johnson AD. Single-nucleotide polymorphism bioinformatics: A comprehensive review of resources. *Circ Cardiovasc Genet* 2009 Oct;2(5):530-6.
92. PHEX gene mutations and genotype-phenotype analysis of korean patients with hypophosphatemic ricketts. EPOS/IFPOS combined meeting; April 11 - 14, 2007; .
93. Kannabiran C, Rogan PK, Olmos L, Basti S, Rao GN, Kaiser-Kupfer M, Hejtmancik JF. Autosomal dominant zonular cataract with sutural opacities is associated with a splice mutation in the betaA3/A1-crystallin gene. *Mol Vis* 1998 Oct 23;4:21.
94. Kaput J, Cotton RG, Hardman L, Watson M, Al Aqeel AI, Al-Aama JY, Al-Mulla F, Alonso S, Aretz S, Auerbach AD, et al. Planning the human variome project: The spain report. *Hum Mutat* 2009 Apr;30(4):496-510.
95. Keren B, Suzuki OT, Gerard-Blanluet M, Bremond-Gignac D, Elmaleh M, Titomanlio L, Delezoide AL, Passos-Bueno MR, Verloes A. CNS malformations in knobloch syndrome with splice mutation in COL18A1 gene. *Am J Med Genet A* 2007 Jul 1;143A(13):1514-8.
96. Kern JS. The molecular basis of dystrophic epidermolysis bullosa: Mutation detection and study of clinical, biochemical, and molecular findings in 29 patients. 2005.
97. A novel intronic point mutation of *CPS1* gene in a korean family with CPS1 deficiency. American society of human genetics meeting; 2007; .

98. Kolsch H, Jessen F, Wiltfang J, Lewczuk P, Dichgans M, Kornhuber J, Frolich L, Heuser I, Peters O, Schulz JB, et al. Influence of SORL1 gene variants: Association with CSF amyloid-beta products in probable alzheimer's disease. *Neurosci Lett* 2008 Jul 25;440(1):68-71.
99. Kolsch H, Jessen F, Wiltfang J, Lewczuk P, Dichgans M, Kornhuber J, Frolich L, Heuser I, Peters O, Schulz JB, et al. Influence of SORL1 gene variants: Association with CSF amyloid-beta products in probable alzheimer's disease. *Neurosci Lett* 2008 Jul 25;440(1):68-71.
100. Kolsch H, Lutjohann D, Jessen F, Popp J, Hentschel F, Kelemen P, Friedrichs S, Maier TA, Heun R. RXRA gene variations influence alzheimer's disease risk and cholesterol metabolism. *J Cell Mol Med* 2009 Mar;13(3):589-98.
101. Kolsch H, Lutjohann D, Jessen F, Popp J, Hentschel F, Kelemen P, Schmitz S, Maier W, Heun R. CYP46A1 variants influence alzheimer's disease risk and brain cholesterol metabolism. *Eur Psychiatry* 2009 Apr;24(3):183-90.
102. Koukouritaki SB, Poch MT, Cabacungan ET, McCarver DG, Hines RN. Discovery of novel flavin-containing monooxygenase 3 (FMO3) single nucleotide polymorphisms and functional analysis of upstream haplotype variants. *Mol Pharmacol* 2005 Aug;68(2):383-92.
103. Kralovicova J, Lei H, Vorechovsky I. Phenotypic consequences of branch point substitutions. *Hum Mutat* 2006 Aug;27(8):803-13.
104. Královicová J, Gaunt TR, Rodriguez S, Wood PJ. Variants in the human insulin gene that affect pre-mRNA splicing: Is -23HphI a functional single nucleotide polymorphism at IDDM2? *Diabetes* 2006 Jan 2006;55(1):260-4.
105. Lacroix M, Lacaze-Buzy L, Furio L, Tron E, Valari M, Van der Wier G, Bodemer C, Bygum A, Bursztejn AC, Gaitanis G, et al. Clinical expression and new SPINK5 splicing defects in netherton syndrome: Unmasking a frequent founder synonymous mutation and unconventional intronic mutations. *J Invest Dermatol* 2011 Nov 17.
106. Lamba V, Lamba J, Yasuda K, Strom S, Davila J, Hancock ML, Fackenthal JD, Rogan PK, Ring B, Wrighton SA, et al. Hepatic CYP2B6 expression: Gender and ethnic differences and relationship to CYP2B6 genotype and CAR (constitutive androstane receptor) expression. *J Pharmacol Exp Ther* 2003 Dec;307(3):906-22.
107. Lancellotti S, Di Leo E, Calandra S, Tarugi P. DIFETTO DI SPLICING DEL PRE-mRNA DELL'APOLIPOPROTEINA B NEL FEGATO DI PAZIENTI CON IPOBETALIPOPROTEINEMIA FAMILIARE. *Patologia Genetica* 2005.
108. Lee PY. Prioritizing SNPs for disease-gene association studies: Algorithms and systems. 2009.
109. Lee PP, Chen TX, Jiang LP, Chen J, Chan KW, Lee TL, Ho MH, Nong SH, Yang Y, Fang YJ, et al. Clinical and molecular characteristics of 35 chinese children with wiskott-aldrich syndrome. *J Clin Immunol* 2009 Jul;29(4):490-500.
110. Lee ST, Lee J, Lee M, Kim JW, Ki CS. Clinical and genetic analysis of korean patients with congenital insensitivity to pain with anhidrosis. *Muscle Nerve* 2009 Nov;40(5):855-9.

111. Leman AR, Pearce DA, Rothberg PG. Gene symbol: CLN3. disease: Juvenile neuronal ceroid lipofuscinosis (batten disease). *Hum Genet* 2005 Feb;116(3):236.
112. Leverenz JB, Yu CE, Montine TJ, Steinbart E, Bekris LM, Zabetian C, Kwong LK, Lee VM, Schellenberg GD, Bird TD. A novel progranulin mutation associated with variable clinical presentation and tau, TDP43 and alpha-synuclein pathology. *Brain* 2007 May;130(Pt 5):1360-74.
113. Li L, Xiao X, Li S, Jia X, Wang P, Guo X, Jiao X, Zhang Q, Hejtmancik JF. Detection of variants in 15 genes in 87 unrelated chinese patients with leber congenital amaurosis. *PLoS One* 2011;6(5):e19458.
114. Lietman SA. Preimplantation genetic diagnosis for hereditary endocrine disease. *Endocr Pract* 2011 Jul-Aug;17 Suppl 3:28-32.
115. Lietman SA, Goldfarb J, Desai N, Levine MA. Preimplantation genetic diagnosis for severe albright hereditary osteodystrophy. *J Clin Endocrinol Metab* 2008 Mar;93(3):901-4.
116. Lim BC, Hwang H, Chae JH, Choi JE, Hwang YS, Kang SH, Ki CS, Kim KJ. SCN1A mutational analysis in korean patients with dravet syndrome. *Seizure* 2011 Dec;20(10):789-94.
117. Lim BC, Ki CS, Kim JW, Cho A, Kim MJ, Hwang H, Kim KJ, Hwang YS, Park WY, Lim YJ, et al. Fukutin mutations in congenital muscular dystrophies with defective glycosylation of dystroglycan in korea. *Neuromuscul Disord* 2010 Aug;20(8):524-30.
118. Liu J, Zhou X, Shan Z, Yang J, Yang Q, Cui Y, Han J. The association of LRP5 gene polymorphisms with ankylosing spondylitis in a chinese han population. *J Rheumatol* 2011 Dec;38(12):2616-8.
119. Liu Z, Venkatesh SS, Maley CC. Sequence space coverage, entropy of genomes and the potential to detect non-human DNA in human samples. *BMC Genomics* 2008 Oct 30;9:509.
120. Lopez-Jimenez E, de Campos JM, Kusak EM, Landa I, Leskela S, Montero-Conde C, Leandro-Garcia LJ, Vallejo LA, Madrigal B, Rodriguez-Antona C, et al. SDHC mutation in an elderly patient without familial antecedents. *Clin Endocrinol (Oxf)* 2008 Dec;69(6):906-10.
121. Lopezjimenez N, Flannick J, Yahyavi M, Li J, Bardakjian T, Tonkin L, Schneider A, Sherr EH, Slavotinek AM. Targeted next-generation sequencing in anophthalmia and microphthalmia patients confirms *SOX2*, *OTX2* and *FOXE3* mutations. *BMC Med Genet* 2011 Dec 28;12(1):172.
122. Luquin N, Yu B, Trent RJ, Morahan JM, Pamphlett R. An analysis of the entire SOD1 gene in sporadic ALS. *Neuromuscul Disord* 2008 Jul;18(7):545-52.
123. Luquin N, Yu B, Saunderson RB, Trent RJ, Pamphlett R. Genetic variants in the promoter of TARDBP in sporadic amyotrophic lateral sclerosis. *Neuromuscular Disorders* 2009 OCT;19(10):696-700.
124. Mackay DS, Henderson RH, Sergouniotis PI, Li Z, Moradi P, Holder GE, Waseem N, Bhattacharya SS, Aldahmesh MA, Alkuraya FS, et al. Novel mutations in MERTK associated with childhood onset rod-cone dystrophy. *Mol Vis* 2010 Mar 9;16:369-77.

125. Maddalena A, Bale S, Das S, Grody W, Richards S, ACMG Laboratory Quality Assurance Committee. Technical standards and guidelines: Molecular genetic testing for ultra-rare disorders. *Genet Med* 2005 Oct;7(8):571-83.
126. Marco EJ, Abidi FE, Bristow J, Dean WB, Cotter P, Jeremy RJ, Schwartz CE, Sherr EH. ARHGEF9 disruption in a female patient is associated with X linked mental retardation and sensory hyperarousal. *BMJ Case Rep* 2009;2009:bcr06.2009.1999. Epub 2009 Jul 2.
127. *ARHGEF9*: Identification of a novel X-linked mental retardation and behavior disorder gene. ; 2006. .
128. Martoni E, Urciuolo A, Sabatelli P, Fabris M, Bovolenta M, Neri M, Grumati P, D'Amico A, Pane M, Mercuri E, et al. Identification and characterization of novel collagen VI non-canonical splicing mutations causing ullrich congenital muscular dystrophy. *Hum Mutat* 2009 May;30(5):E662-72.
129. Maruszak A, Safranow K, Gustaw K, Kijanowska-Haladyna B, Jakubowska K, Olszewska M, Styczynska M, Berdyski M, Tysarowski A, Chlubek D, et al. PIN1 gene variants in alzheimer's disease. *BMC Med Genet* 2009 Nov 12;10:115.
130. Megremis S, Mitsioni A, Mitsioni AG, Fylaktou I, Kitsiou-Tzelli S, Stefanidis CJ, Kanavakis E, Traeger-Synodinos J. Nucleotide variations in the NPHS2 gene in greek children with steroid-resistant nephrotic syndrome. *Genet Test Mol Biomarkers* 2009 Apr;13(2):249-56.
131. Mintchev N, Zamba-Papanicolaou E, Kleopa KA, Christodoulou K. A novel ALS2 splice-site mutation in a cypriot juvenile-onset primary lateral sclerosis family. *Neurology* 2009 Jan 6;72(1):28-32.
132. Genotype and tissue effects on alternative splicing of the TCF7L2 gene in tissues important to type 2 diabetes (T2DM) pathogenesis. Abstract book #1352; November 11-15, 2008; The American Society of Human Genetics.
133. Moriwaki K, Noda K, Furukawa Y, Ohshima K, Uchiyama A, Nakagawa T, Taniguchi N, Daigo Y, Nakamura Y, Hayashi N, et al. Deficiency of GMDS leads to escape from NK cell-mediated tumor surveillance through modulation of TRAIL signaling. *Gastroenterology* 2009 Jul;137(1):188,98, 198.e1-2.
134. Mucaki EJ, Ainsworth P, Rogan PK. Comprehensive prediction of mRNA splicing effects of BRCA1 and BRCA2 variants. *Hum Mutat* 2011 Jul;32(7):735-42.
135. Mukhopadhyay A, Nikopoulos K, Maugeri A, de Brouwer AP, van Nouhuys CE, Boon CJ, Perveen R, Zegers HA, Wittebol-Post D, van den Biesen PR, et al. Erosive vitreoretinopathy and wagner disease are caused by intronic mutations in CSPG2/Versican that result in an imbalance of splice variants. *Invest Ophthalmol Vis Sci* 2006 Aug;47(8):3565-72.
136. Najah M, Di Leo E, Awatef J, Magnolo L, Imene J, Pinotti E, Bahri M, Barsaoui S, Brini I, Fekih M, et al. Identification of patients with abetalipoproteinemia and homozygous familial hypobetalipoproteinemia in tunisia. *Clin Chim Acta* 2009 Mar;401(1-2):51-6.
137. Nalla VK. Automated splice site analysis. 2005.
138. Nalla VK, Rogan PK. Automated splicing mutation analysis by information theory. *Hum Mutat* 2005 Apr;25(4):334-42.

139. Nalla VK, Rogan PK. Automated splicing mutation analysis by information theory (vol 25, pg 334, 2005). *Hum Mutat* 2008 SEP;29(9):1168-.
140. Naruse H, Ikawa N, Yamaguchi K, Nakamura Y, Arai M, Ishioka C, Sugano K, Tamura K, Tomita N, Matsubara N, et al. Determination of splice-site mutations in lynch syndrome (hereditary non-polyposis colorectal cancer) patients using functional splicing assay. *Fam Cancer* 2009;8(4):509-17.
141. Nasim MT, Ogo T, Ahmed M, Randall R, Chowdhury HM, Snape KM, Bradshaw TY, Southgate L, Lee GJ, Jackson I, et al. Molecular genetic characterization of SMAD signaling molecules in pulmonary arterial hypertension. *Hum Mutat* 2011 DEC;32(12):1385-9.
142. Nasim MT, Ogo T, Ahmed M, Randall R, Chowdhury HM, Snape KM, Bradshaw TY, Southgate L, Lee GJ, Jackson I, et al. Molecular genetic characterization of SMAD signaling molecules in pulmonary arterial hypertension. *Hum Mutat* 2011 DEC;32(12):1385-9.
143. Oetting WS, Tabone T. The 2004 human genome variation society scientific meeting. *Hum Mutat* 2005 Aug;26(2):160-3.
144. Oh KS, Khan SG, Jaspers NG, Raams A, Ueda T, Lehmann A, Friedmann PS, Emmert S, Gratchev A, Lachlan K, et al. Phenotypic heterogeneity in the XPB DNA helicase gene (ERCC3): Xeroderma pigmentosum without and with cockayne syndrome. *Hum Mutat* 2006 Nov;27(11):1092-103.
145. Oh SW, Lee JS, Kim MY, Kim SC. COL7A1 mutational analysis in korean patients with dystrophic epidermolysis bullosa. *Br J Dermatol* 2007 Dec;157(6):1260-4.
146. Oh SW, Lee JS, Kim MY, Kim SC. Novel keratin 5 mutations in epidermolysis bullosa simplex: Cases with unusual genotype-phenotype correlation. *J Dermatol Sci* 2007 Dec;48(3):229-32.
147. Okubo M, Ishihara M, Iwasaki T, Ebara T, Aoyama Y, Murase T, Hattori H. A novel APOA5 splicing mutation IVS2+1g > a in a japanese chylomicronemia patient. *Atherosclerosis* 2009 NOV;207(1):24-5.
148. O'Neill JP, Rogan PK, Cariello N, Nicklas JA. Mutations that alter RNA splicing of the human HPRT gene: A review of the spectrum. *Mutat Res* 1998 Nov;411(3):179-214.
149. Ozaltin F, Ibsirlioglu T, Taskiran EZ, Baydar DE, Kaymaz F, Buyukcelik M, Kilic BD, Balat A, Iatropoulos P, Asan E, et al. Disruption of PTPRO causes childhood-onset nephrotic syndrome. *Am J Hum Genet* 2011 Jul 15;89(1):139-47.
150. Palomino-Doza J, Rahman TJ, Avery PJ, Mayosi BM, Farrall M, Watkins H, Edwards CR, Keavney B. Ambulatory blood pressure is associated with polymorphic variation in P2X receptor genes. *Hypertension* 2008 Nov;52(5):980-5.
151. Papi L, Putignano AL, Congregati C, Piaceri I, Zanna I, Sera F, Morrone D, Genuardi M, Palli D. A PALB2 germline mutation associated with hereditary breast cancer in italy. *Fam Cancer* 2010 Jun;9(2):181-5.
152. Papi L, Putignano AL, Congregati C, Piaceri I, Zanna I, Sera F, Morrone D, Genuardi M, Palli D. A PALB2 germline mutation associated with hereditary breast cancer in italy RID C-8176-2011. *Familial Cancer* 2010 JUN;9(2):181-5.

153. Papp J, Kovacs ME, Olah E. Germline MLH1 and MSH2 mutational spectrum including frequent large genomic aberrations in hungarian hereditary non-polyposis colorectal cancer families: Implications for genetic testing. *World J Gastroenterol* 2007 May 21;13(19):2727-32.
154. Parslow GR. Websites of note. *Biochem Mol Biol Educ* 2011 May-Jun;39(3):230-2.
155. Pasmooij AM. Revertant mosaicism in epidermolysis bullosa due to different second site mutations in LAMB3. 2006:Chapter 3.
156. Pasmooij AM, Pas HH, Bolling MC, Jonkman MF. Revertant mosaicism in junctional epidermolysis bullosa due to multiple correcting second-site mutations in LAMB3. *J Clin Invest* 2007 May;117(5):1240-8.
157. Pasvolsky R, Feigelson SW, Kilic SS, Simon AJ, Tal-Lapidot G, Grabovsky V, Crittenden JR, Amariglio N, Safran M, Graybiel AM, et al. A LAD-III syndrome is associated with defective expression of the rap-1 activator CalDAG-GEFI in lymphocytes, neutrophils, and platelets. *J Exp Med* 2007 Jul 9;204(7):1571-82.
158. Pasvolsky R, Feigelson SW, Kilic SS, Simon AJ, Tal-Lapidot G, Grabovsky V, Crittenden JR, Amariglio N, Safran M, Graybiel AM, et al. A LAD-III syndrome is associated with defective expression of the rap-1 activator CalDAG-GEFI in lymphocytes, neutrophils, and platelets. *J Exp Med* 2007 JUL 9;204(7):1571-82.
159. Pelucchi S, Mariani R, Trombini P, Coletti S, Pozzi M, Paolini V, Barisani D, Piperno A. Expression of hepcidin and other iron-related genes in type 3 hemochromatosis due to a novel mutation in transferrin receptor-2. *Haematologica* 2009 Feb;94(2):276-9.
160. Piva F, Giulietti M, Nardi B, Bellantuono C, Principato G. An improved in silico selection of phenotype affecting polymorphisms in SLC6A4, HTR1A and HTR2A genes. *Human Psychopharmacology-Clinical and Experimental* 2010 MAR;25(2):153-61.
161. Priore Oliva C, Tarugi P, Calandra S, Pisciotta L, Bellocchio A, Bertolini S, Guardamagna O, Schaap FG. A novel sequence variant in APOA5 gene found in patients with severe hypertriglyceridemia. *Atherosclerosis* 2006 Sep;188(1):215-7.
162. Qin S, Shen L, Zhang A, Xie J, Shen W, Chen L, Tang J, Xiong Y, Yang L, Shi Y, et al. Systematic polymorphism analysis of the CYP2D6 gene in four different geographical han populations in mainland china. *Genomics* 2008 Sep;92(3):152-8.
163. Rhyne J, Mantaring MM, Gardner DF, Miller M. Multiple splice defects in ABCA1 cause low HDL-C in a family with hypoalphalipoproteinemia and premature coronary disease. *BMC Med Genet* 2009 Jan 8;10:1.
164. Riveira-Munoz E, Chang Q, Godefroid N, Hoenderop JG, Bindels RJ, Dahan K, Devuyst O, Belgian Network for Study of Gitelman Syndrome. Transcriptional and functional analyses of SLC12A3 mutations: New clues for the pathogenesis of gitelman syndrome. *J Am Soc Nephrol* 2007 Apr;18(4):1271-83.
165. Riveira-Munoz E, Devuyst O, Belge H, Jeck N, Strompf L, Vargas-Poussou R, Jeunemaitre X, Blanchard A, Knoers NV, Konrad M, et al. Evaluating PVALB as a candidate gene for SLC12A3-negative cases of gitelman's syndrome. *Nephrol Dial Transplant* 2008 Oct;23(10):3120-5.

166. Rogan PK, Faux BM, Schneider TD. Information analysis of human splice site mutations. *Hum Mutat* 1998;12(3):153-71.
167. Rogan PK, Schneider TD. Using information content and base frequencies to distinguish mutations from genetic polymorphisms in splice junction recognition sites. *Hum Mutat* 1995;6(1):74-6.
168. Rogan PK, Svojanovsky S, Leeder JS. Information theory-based analysis of CYP2C19, CYP2D6 and CYP3A5 splicing mutations. *Pharmacogenetics* 2003 Apr;13(4):207-18.
169. Rogan PK. Ab initio exon definition using an information theory-based approach. ; 2009. PT: B; CT: 43rd Annual Conference on Information Sciences and Systems; CY: MAR 18-20, 2009; CL: Baltimore, MD; TC: 0; UT: WOS:000270992700158.
170. Rossi PI, Vaccari CM, Terracciano A, Doria-Lamba L, Facchinetti S, Priolo M, Ayuso C, De Jorge L, Gimelli S, Santorelli FM, et al. The metabotropic glutamate receptor 1, GRM1: Evaluation as a candidate gene for inherited forms of cerebellar ataxia. *J Neurol* 2010 Apr;257(4):598-602.
171. Rossi PI, Vaccari CM, Terracciano A, Doria-Lamba L, Facchinetti S, Priolo M, Ayuso C, De Jorge L, Gimelli S, Santorelli FM, et al. The metabotropic glutamate receptor 1, GRM1: Evaluation as a candidate gene for inherited forms of cerebellar ataxia. *J Neurol* 2010 Apr;257(4):598-602.
172. Roux-Buisson N, Rendu J, Denjoy I, Guicheney P, Goldenberg A, David N, Faivre L, Barthez O, Danieli GA, Marty I, et al. Functional analysis reveals splicing mutations of the CASQ2 gene in patients with CPVT: Implication for genetic counselling and clinical management. *Hum Mutat* 2011 May 26.
173. Russcher H. Glucocorticoid receptor variants modulate the sensitivity to cortisol. 2006.
174. Russcher H, Smit P, van Rossum EF, van den Akker EL, Brinkmann AO, de Heide LJ, de Jong FH, Koper JW, Lamberts SW. Strategies for the characterization of disorders in cortisol sensitivity. *J Clin Endocrinol Metab* 2006 Feb;91(2):694-701.
175. Sabet A, Li J, Ghandour K, Pu Q, Wu X, Kamholz J, Shy ME, Cambi F. Skin biopsies demonstrate MPZ splicing abnormalities in charcot-marie-tooth neuropathy 1B. *Neurology* 2006 Oct 10;67(7):1141-6.
176. Sanggaard KM, Rendtorff ND, Kjaer KW, Eiberg H, Johnsen T, Gimsing S, Dyrmoose J, Nielsen KO, Lage K, Tranebjaerg L. Branchio-oto-renal syndrome: Detection of EYA1 and SIX1 mutations in five out of six danish families by combining linkage, MLPA and sequencing analyses. *Eur J Hum Genet* 2007 Nov;15(11):1121-31.
177. Schneider TD, Rogan PK, inventors. Computational analysis of nucleic acid information defines binding sites. The United States of America as represented by the Department of Health and Human Services, editor. USA patent 5867402 Feb 2, 1999.
178. Schonfelder EM, Knuppel T, Tasic V, Miljkovic P, Konrad M, Wuhl E, Antignac C, Bakkaloglu A, Schaefer F, Weber S, et al. Mutations in uroplakin IIIA are a rare cause of renal hypodysplasia in humans. *Am J Kidney Dis* 2006 Jun;47(6):1004-12.
179. Schwaderer P, Knuppel T, Konrad M, Mehls O, Scharer K, Schaefer F, Weber S. Clinical course and NPHS2 analysis in patients with late steroid-resistant nephrotic syndrome. *Pediatr Nephrol* 2008 Feb;23(2):251-6.

180. Simpson MA, Hsu R, Keir LS, Hao J, Sivapalan G, Ernst LM, Zackai EH, Al-Gazali LI, Hulskamp G, Kingston HM, et al. Mutations in FAM20C are associated with lethal osteosclerotic bone dysplasia (raine syndrome), highlighting a crucial molecule in bone development. *Am J Hum Genet* 2007 Nov;81(5):906-12.
181. Skipper L, Shen H, Chua E, Bonnard C, Kolatkar P, Tan LC, Jamora RD, Puvan K, Puong KY, Zhao Y, et al. Analysis of LRRK2 functional domains in nondominant parkinson disease. *Neurology* 2005 Oct 25;65(8):1319-21.
182. Slavotinek AM, Baranzini SE, Schanze D, Labelle-Dumais C, Short KM, Chao R, Yahyavi M, Bijlsma EK, Chu C, Musone S, et al. Manitoba-oculo-tricho-anal (MOTA) syndrome is caused by mutations in FREM1. *J Med Genet* 2011 Jun;48(6):375-82.
183. Slavotinek AM, Chao R, Vacik T, Yahyavi M, Abouzeid H, Bardakjian T, Schneider A, Shaw G, Sherr EH, Lemke G, et al. VAX1 mutation associated with microphthalmia, corpus callosum agenesis, and orofacial clefting: The first description of a VAX1 phenotype in humans. *Hum Mutat* 2012 Feb;33(2):364-8.
184. Smit P. Factors determining glucocorticoid sensitivity in man. 2006.
185. Song HR, Park JW, Cho DY, Yang JH, Yoon HR, Jung SC. PHEX gene mutations and genotype-phenotype analysis of korean patients with hypophosphatemic rickets. *J Korean Med Sci* 2007 Dec;22(6):981-6.
186. Soran H, Charlton-Menys V, Hegele R, Wang J, Benbow EW, Roberts I, Wood G, Durrington P. Proteinuria and severe mixed dyslipidemia associated with a novel APOAV gene mutation. *J Clin Lipidol* 2010 Jul-Aug;4(4):310-3.
187. Spurdle AB, Couch FJ, Hogervorst FB, Radice P, Sinilnikova OM, IARC Unclassified Genetic Variants Working Group. Prediction and assessment of splicing alterations: Implications for clinical testing. *Hum Mutat* 2008 Nov;29(11):1304-13.
188. Stockley TL, Mendoza-London R, Propst EJ, Sodhi S, Dupuis L, Papsin BC. A recurrent EYA1 mutation causing alternative RNA splicing in branchio-oto-renal syndrome: Implications for molecular diagnostics and disease mechanism. *American Journal of Medical Genetics Part a* 2009 MAR;149A(3):322-7.
189. Svojanovsky SR, Schneider TD, Rogan PK. Redundant designations of BRCA1 intron 11 splicing mutation; c. 4216-2A>G; IVS11-2A>G; L78833, 37698, A>G. *Hum Mutat* 2000 Sep;16(3):264.
190. Sznajder Y, Coldea C, Meire F, Delpierre I, Sekhara T, Touraine RL. A de novo SOX10 mutation causing severe type 4 waardenburg syndrome without hirschsprung disease. *Am J Med Genet A* 2008 Apr 15;146A(8):1038-41.
191. Tartaglia-Polcini A, Bonnard C, Micheloni A, Cianfarani F, Andre A, Zambruno G, Hovnanian A, D'Alessio M. SPINK5, the defective gene in netherton syndrome, encodes multiple LEKTI isoforms derived from alternative pre-mRNA processing. *J Invest Dermatol* 2006 Feb;126(2):315-24.
192. Tazi J, Durand S, Jeanteur P. The spliceosome: A novel multi-faceted target for therapy. *Trends Biochem Sci* 2005 Aug;30(8):469-78.
193. The American College of Medical Genetics Laboratory Quality Assurance Committee. Technical standards and guidelines: Molecular genetic testing for rare disorders. ; 2005.

194. Thompson TE, Rogan PK, Risinger JI, Taylor JA. Splice variants but not mutations of DNA polymerase beta are common in bladder cancer. *Cancer Res* 2002 Jun 1;62(11):3251-6.
195. Titeux M, Mejia JE, Mejlumian L, Bourthoumieu S, Mirval S, Tonasso L, Heller M, Prost-Squarcioni C, Hovnanian A. Recessive dystrophic epidermolysis bullosa caused by COL7A1 hemizygoty and a missense mutation with complex effects on splicing. *Hum Mutat* 2006 Mar;27(3):291-2.
196. Glial cell-line derived neurotrophic factor: Un gene candidate per la patogenesi del rene con midollare a spugna [Internet]; c2006. Available from: http://www.sigu.net/e107_files/downloads/Comunicazioni/Genetica%20delle%20Malattie%20Complesse.pdf.
197. Tosetto E, Ceol M, Mezzabotta F, Ammenti A, Peruzzi L, Caruso MR, Barbano G, Vezzoli G, Colussi G, Vergine G, et al. Novel mutations of the CLCN5 gene including a complex allele and A 5' UTR mutation in dent disease 1. *Clin Genet* 2009 Oct;76(4):413-6.
198. Tosetto E, Ghiggeri GM, Emma F, Barbano G, Carrea A, Vezzoli G, Torregrossa R, Cara M, Ripanti G, Ammenti A, et al. Phenotypic and genetic heterogeneity in dent's disease--the results of an italian collaborative study. *Nephrol Dial Transplant* 2006 Sep;21(9):2452-63.
199. Tournier I, Vezain M, Martins A, Charbonnier F, Baert-Desurmont S, Olschwang S, Wang Q, Buisine MP, Soret J, Tazi J, et al. A large fraction of unclassified variants of the mismatch repair genes MLH1 and MSH2 is associated with splicing defects. *Hum Mutat* 2008 Dec;29(12):1412-24.
200. Tram E, Ibrahim-Zada I, Briollais L, Knight JA, Andrulis IL, Ozcelik H. Identification of germline alterations of the mad homology 2 domain of SMAD3 and SMAD4 from the ontario site of the breast cancer family registry (CFR). *Breast Cancer Res* 2011 Aug 11;13(4):R77.
201. Tunca B, Pedroni M, Cecener G, Egeli U, Borsi E, Zorluoglu A, Di Gregorio C, Yilmazlar T, Yerci O, de Leon MP. Analysis of mismatch repair gene mutations in turkish HNPCC patients. *Fam Cancer* 2010 Sep;9(3):365-76.
202. Tuohy T, Burt RW. Attenuated familial adenomatous polyposis: Diagnosis, management and future prognosis. In: R. Rodriguez-Bigas, R. Cutait, P. Lunch, I. Tomlinson, H. Vasen, editors. *Hereditary colorectal cancer*. Springer; 2010. .
203. Vockley J, Rogan PK, Anderson BD, Willard J, Seelan RS, Smith DI, Liu W. Exon skipping in IVD RNA processing in isovaleric acidemia caused by point mutations in the coding region of the IVD gene. *Am J Hum Genet* 2000 Feb;66(2):356-67.
204. von Kodolitsch Y, Berger J, Rogan PK. Predicting severity of haemophilia A and B splicing mutations by information analysis. *Haemophilia* 2006 May;12(3):258-62.
205. von Kodolitsch Y, Berger J, Rogan PK. Predicting severity of haemophilia A and B splicing mutations by information analysis. *Haemophilia* 2006 May;12(3):258-62.
206. von Kodolitsch Y, Nienaber CA, Fliegner M, Rogan PK. Splice site mutations and atherosclerosis: Mechanisms and prediction models. *Z Kardiol* 2001 Feb;90(2):87-95.
207. von Kodolitsch Y, Pyeritz RE, Rogan PK. Splice-site mutations in atherosclerosis candidate genes: Relating individual information to phenotype. *Circulation* 1999 Aug 17;100(7):693-9.

208. Vorechovsky I. Aberrant 3' splice sites in human disease genes: Mutation pattern, nucleotide structure and comparison of computational tools that predict their utilization. *Nucleic Acids Res* 2006;34(16):4630-41.
209. Vreeswijk MP, Kraan JN, van der Klift HM, Vink GR, Cornelisse CJ, Wijnen JT, Bakker E, van Asperen CJ, Devilee P. Intronic variants in BRCA1 and BRCA2 that affect RNA splicing can be reliably selected by splice-site prediction programs. *Hum Mutat* 2009 Jan;30(1):107-14.
210. Wan L, Lee CC, Hsu CM, Hwu WL, Yang CC, Tsai CH, Tsai FJ. Identification of eight novel mutations of the acid alpha-glucosidase gene causing the infantile or juvenile form of glycogen storage disease type II. *J Neurol* 2008 Jun;255(6):831-8.
211. Wang E, Dimova N, Cambi F. PLP/DM20 ratio is regulated by hnRNPH and F and a novel G-rich enhancer in oligodendrocytes. *Nucleic Acids Res* 2007;35(12):4164-78.
212. Wang J, Sonnerborg A, Rane A, Josephson F, Lundgren S, Stahle L, Ingelman-Sundberg M. Identification of a novel specific CYP2B6 allele in africans causing impaired metabolism of the HIV drug efavirenz. *Pharmacogenet Genomics* 2006 Mar;16(3):191-8.
213. Wang P, Guo X, Jia X, Li S, Xiao X, Zhang Q. Novel mutations of the PAX6 gene identified in chinese patients with aniridia. *Mol Vis* 2006 Jun 7;12:644-8.
214. Watnick TJ, Garcia-Gonzalez MA, Germino GG, Jones JG, inventors. PKD mutations and evaluation of the same. International patent . 2008.
215. Wessagowit V, Kim SC, Woong Oh S, McGrath JA. Genotype-phenotype correlation in recessive dystrophic epidermolysis bullosa: When missense doesn't make sense. *J Invest Dermatol* 2005 Apr;124(4):863-6.
216. Wessagowit V, McGrath JA. Clinical and molecular significance of splice site mutations in the plakophilin 1 gene in patients with ectodermal dysplasia-skin fragility syndrome. *Acta Derm Venereol* 2005;85(5):386-8.
217. Wessagowit V, Nalla VK, Rogan PK, McGrath JA. Normal and abnormal mechanisms of gene splicing and relevance to inherited skin diseases. *J Dermatol Sci* 2005 Nov;40(2):73-84.
218. Modeling aberrant splicing in mutant genes associated with inherited retinal degeneration and bardet-biedl syndrome. *American society of human genetics 2008 - annual meeting; 2008; American Society of Human Genetics 2008.*
219. Wong T, Gammon L, Liu L, Mellerio JE, Dopping-Hepenstal PJ, Pacy J, Elia G, Jeffery R, Leigh IM, Navsaria H, et al. Potential of fibroblast cell therapy for recessive dystrophic epidermolysis bullosa. *J Invest Dermatol* 2008 Sep;128(9):2179-89.
220. Xiong Y, Wang M, Fang K, Xing Q, Feng G, Shen L, He L, Qin S. A systematic genetic polymorphism analysis of the CYP2C9 gene in four different geographical han populations in mainland china. *Genomics* 2011 May;97(5):277-81.
221. Xu X, Li S, Xiao X, Wang P, Guo X, Zhang Q. Sequence variations of GRM6 in patients with high myopia. *Mol Vis* 2009 Oct 19;15:2094-100.
222. Yang M, Solidar A, Wyckoff GJ. Novel method for discerning the action of selection during evolution. *J Biomed Sci Eng* 2010;3:109-113.

223. Yu B. Role of in silico tools in gene discovery. *Mol Biotechnol* 2009 Mar;41(3):296-306.
224. Yu H, Patel SB. Recent insights into the smith-lemli-opitz syndrome. *Clin Genet* 2005 Nov;68(5):383-91.
225. Zaffanello M, Taranta A, Palma A, Bettinelli A, Marseglia GL, Emma F. Type IV bartter syndrome: Report of two new cases. *Pediatr Nephrol* 2006 Jun;21(6):766-70.
226. Zampieri S, Buratti E, Dominissini S, Montalvo AL, Pittis MG, Bembi B, Dardis A. Splicing mutations in glycogen-storage disease type II: Evaluation of the full spectrum of mutations and their relation to patients' phenotypes. *Eur J Hum Genet* 2011 Apr;19(4):422-31.
227. Zhang K, Nowak I, Rushlow D, Gallie BL, Lohmann DR. Patterns of missplicing caused by RB1 gene mutations in patients with retinoblastoma and association with phenotypic expression. *Hum Mutat* 2008 Apr;29(4):475-84.
228. Zhang Q, Zulfiqar F, Riazuddin SA, Xiao X, Yasmeen A, Rogan PK, Caruso R, Sieving PA, Riazuddin S, Hejtmancik JF. A variant form of oguchi disease mapped to 13q34 associated with partial deletion of GRK1 gene. *Mol Vis* 2005 Nov 14;11:977-85.